Neuroscience Letters, 121 (1991) 275–278
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Aceves, J., Floran, B., Martinez-Fong, D., Sierra, A., Hernandez, S. and Mariscal, S., L-DOPA stimulates the release of [^{3}H]γ-aminobutyric acid in the basal ganglia of 6-hydroxydopamine lesioned rats, 223
Aguayo, L.G., Weight, F.F. and White, G., TTX-sensitive action potentials and excitability of adult rat sensory neurons cultured in serum- and exogenous nerve growth factor-free medium, 88
Akaike, N., see Yamasaki, Y., 251
Akiyama, H., see Tooyama, I., 155
Allan, R., see Fletcher, E., 133
Allen, S.J., see Treanor, J.J.S., 73
Allsop, D., see Sato, M., 271
Ansselin, A.D., see Davey, D.F., 83
Attanasio, A., see Migheli, A., 55
Aydar, E., see Dickenson, A.H., 263

Bando, Y., see Horie, H., 125
Ban, H., see Yamasaki, Y., 251
Baron, J.C., see Blin, J., 183
Baruah, S., see Sherman, A.D., 77
Behnisch, T., see Matthies Jr., H. 259
Belan, I., see Firth, B.T., 139
Bernard, G., see Lefauconnier, J.M., 9
Bizzini, B., see Chapron, J., 21
Blazynski, C., The accumulation of [^{3}H]phenylisopropyl adenosine ([^{3}H]PIA) and [^{3}H]adenosine into rabbit retinal neurons is inhibited by nitrobenzylthioinosine (NBI), 1
Blin, J., Denis, A., Yamaguchi, T., Crouzel, C., MacKenzie, E.T. and Baron, J.C., PET studies of [^{18}F]methyl-MK-801, a potential NMDA receptor complex radioligand, 183
Block, F., see Wagemann, E., 102
Bolden, S.W., see Rogers, L.J., 63
Borbély, A.A., see Dijk, D.-J., 59
Bouchaud, C., see Lefauconnier, J.M., 9
Bowie, D. and Smart, T.G., Interaction of 5-bromowillardiine with non-NMDA receptors expressed in *Xenopus laevis* oocytes injected with chick brain mRNA, 68
Bubel, H.C., see Keller, J.T., 247
Budd, G.C., see Das, A., 231
Budelmann, B.U., see Williamson, R., 215
Burešová, M., Dvořáková, M., Zvolský, P. and Illnerová, H., Early morning bright light phase advances the human circadian pacemaker within one day, 47

Cajochen, C., see Dijk, D.-J., 59
Carlton, S.M. and Hayes, E.S., GABAergic vesicle-containing dendrites and spines: a critical element in processing sensory input in the monkey dorsal horn, 40
Chan, J. and Greenberg, D.A., Effects of glucose on calcium channels in neural cells, 34
Chapron, J., Vaillier, J., Koenig, J., De la Porte, S. and Bizzini, B., The effect of tetanus toxin on in vitro synaptogenesis, 21
Chase, T.N., see Susel, Z., 234
Chi, H., see Horie, H., 125
Copray, J.C.V.M., Liem, R.S.B., Ter Horst, G.J. and Van Willigen, J.D., Origin, distribution and morphology of serotonergic afferents to the mesencephalic trigeminal nucleus of the rat, 97
Cotman, C., see Peterson, C., 239
Crouzel, C., see Blin, J., 183
Crow, T., Forrester J., Williams M., Waxham, M.N. and Neary, J., Down-regulation of protein kinase C blocks 5-HT-induced enhancement in *Hermissenda* B photoreceptors, 107

Daly, J.W., see Karlsten, R., 267
Darlington, C.L., see Smith, P.F., 244
Das, A., Pansky, B., Budd, G.C. and Reid, T.W., Human retinoblastoma Y79 cells contain both insulin-specific mRNA and insulin-binding sites, 231
Davey, D.F. and Ansselin, A.D., Labelling of restricted numbers of axons by solid rhodamine implantation into nerve trunks, 83
Davidson, A.T., see Sherman, A.D., 77
Dawbarn, D., see Treanor, J.J.S., 73
De la Porte, S., see Chapron, J., 21
Denis, A., see Blin, J., 183
De Reviers, M.-M., Tillet, Y. and Pelletier, J., Melatonin binding sites in the brain of sheep exposed to light or pinealectomized, 17
Dickenson, A.H. and Aydar, E., Antagonism at the glycine site on the NMDA receptor reduces spinal nociception in the rat, 263
Dinopoulos, A., Dori, I. and Parnavelas, J.G., Immunohistochemical localization of aspartate in corticofugal pathways, 25
Dori, I., see Dinopoulos, A., 25
Drew, C., see Fletcher, E., 133
Duron, B., see Larnicol, N., 93
Dvořáková, M., see Burešová, M., 47
Dijk, D.-J., Cajochen, C. and Borbély, A.A., Effect of a single 3-hour exposure to bright light on core body temperature and sleep in humans, 59

Edvinsson, L., Ekman, R., Jansen, I., McCulloch, J., Mortensen, A. and Uddman, R., Reduced levels of calcitonin gene-related peptide-like immunoreactivity in human brain vessels after subarachnoid haemorrhage, 151
Ekman, R., see Edvinsson, L., 151
Engber, T.M., see Susel, Z., 234

Firth, B.T., Kennaway, D.J. and Belan, I., Thermoperiodic influences on plasma melatonin rhythms in the lizard *Tiliqua rugosa*: effect of thermophase duration, 139
Fletcher, E., Mewett, K., Drew, C., Allan, R. and Johnston, G.R., Inhibition of high affinity L-glutamic acid uptake into rat cortical synaptosomes by the conformationally restricted analogue of glutamic acid, *cis*-1-aminocyclobutane-1,3-dicarboxylic acid, 133
Floran, B., see Aceves, J., 223
Forrester, J., see Crow, T., 107
Fukuda, M., see Ono, T., 194

Greenberg, D.A., see Chan, J., 34
Guilarte, T.R., Abnormal endogenous amino acid release in brain slices from vitamin B-6 restricted neonatal rats, 203
Guilarte, T.R., Reduced NMDA receptor-ion channel function in the vitamin B-6 restricted neonatal rat brain., 207
Guy, S.P., Jones, D., Mann, D.M.A. and Itzhaki, R.F., Human neuroblastoma cells treated with aluminium express an epitope associated with Alzheimer's disease neurofibrillary tangles, 166

Haga, S., see Sato, M., 271
Hara, H., see Yamasaki, Y., 251
Hara, Ya., see Tooyama, I., 155

Hayakawa, T., see Kohmura, E., 159
Hayes, E.S., see Carlton, S.M., 40
Hegwood, T.S., see Sherman, A.D., 77
Henry, M.A., see Westrum, L.E., 143
Hernandez, S., see Aceves, J., 223
Hidaka, H., see Ito, I., 119
Hide, I., see Karlsten, R., 267
Horie, H., Bando, Y., Chi, H. and Takenaka, T., NGF enhances neurite regeneration from nerve-transected terminals of young adult and aged mouse dorsal root ganglia in vitro, 125
Houchi, H., see Nakanishi, A., 163
Hubbard, J.I., see Smith, P.F., 244

Ichikawa, Y., Terakado, Y. and Yamaguchi, T., Last-order interneurones controlling activity of elbow extensor motoneurones during forelimb fictive locomotion in the cat, 37
Ikeda, K., see Sato, M., 271
Illnerová, H., see Burešová, M., 47
Ishii, T., see Sato, M., 271
Ito, I., Hidaka, H. and Sugiyama, H., Effects of KN-62, a specific inhibitor of calcium/calmodulin-dependent protein kinase II, on long-term potentiation in the rat hippocampus, 119

Jaarsma, D., Sebens, J. and Korf, J., Reduction of adenosine A_1-receptors in the perforant pathway terminal zone in Alzheimer hippocampus, 111
Jansen, I., see Edvinsson, L., 151
Jarolimek, W. and Misgeld, U., Reduction of $GABA_A$ receptor-mediated inhibition by the non-NMDA receptor antagonist 6-cyano-7-nitroquinoxaline-2,3-dione in cultured neurons of rat brain, 227
Jen, L.S., see Lau, K.C., 255
Johnson, K.M., see Sacaan, A.I., 219
Johnston, G.R., see Fletcher, E., 133
Jones, D., see Guy, S.P., 166

Karlsten, R., Post, C., Hide, I. and Daly, J.W., The antinociceptive effect of intrathecally administered adenosine analogs in mice correlates with the affinity for the A_1-adenosine receptor, 267
Kase, H., see Matthies Jr., H. 259
Keller, J.T., Bubel, H.C., Wander, A.H. and Tierney, B.E., Somatotopic distribution of corneal afferent neurons in the guinea pig trigeminal ganglion, 247
Kennaway, D.J., see Firth, B.T., 139
Kerwin, J., Morris, C., Oakley, A., Perry, R. and Perry, E., Distribution of nerve growth factor receptor immunoreactivity in the human hippocampus, 178
Kimura, H., see Tooyama, I., 155
Kinoshita, A., see Kohmura, E., 159
Koenig, J., see Chapron, J., 21
Kogure, K., see Yamasaki, Y., 251
Kohmura, E., Yamada, K., Hayakawa, T. and Kinoshita, A., Neurotoxicity caused by glutamate after subcritical hypoxia is prevented by 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX): an in vitro study using rat hippocampal neurons, 159
Koob, G.F., see Schroeder, R.L., 173
Kort, J., see Jaarsma, D., 111
Koyano, K., see Tanaka, K., 191
Kuba, K., see Tanaka, K., 191
Kuo, S., see Susel, Z., 234
Kurogochi, Y., see Nakayama, H., 122

Lambert, G.A., see Zagami, A.S., 187
Larnicol, N., Rose, D. and Duron, B., Postnatal development of small and large dorsal root ganglion neurons in the cat. A study on cervical levels (C_5–C_6) and on phrenic afferents, 93
Lau, K.C., So, K. -F., Tay, D. and Jen, L.S., Elimination of transient dendritic spines in ipsilaterally projecting retinal ganglion cells in rats with neonatal unilateral thalamotomy, 255
Lefauconnier, J.M., Bouchaud, C. and Bernard, G., Initial process of diffusion of small molecules from blood vessels to the meninges in the young rat, 9
Lehmenkühler, C., Walden, J. and Speckmann, E. -J., Decrease of *N*-methyl-D-aspartate responses by noradrenaline in the rat motorcortex in vivo, 5
Liem, R.S.B., see Copray, J.C.V.M., 97
Linnoila, M., see Ota, M., 129
Lynch, G., see Peterson, C., 239

MacGowan, S.H., see Treanor, J.J.S., 73
MacKenzie, E.T., see Blin, J., 183
Mahata, S., Effect of insulin on serotonin, 5-hydroxyindole-acetic acid, norepinephrine, epinephrine and corticosterone contents in chick, 115
Mann, D.M.A., see Guy, S.P., 166
Mariscal, S., see Aceves, J., 223
Martinez-Fong, D., see Aceves, J., 223
Matthies Jr., H. Behnisch, T., Kase, H., Matthies, H. and Reymann, K.G., Differential effects of protein kinase inhibitors on pre-established long-term potentiation in rat hippocampal neurons in vitro, 259
Matthies, H., see Matthies Jr., H. 259
McCulloch, J., see Edvinsson, L., 151
McGeer, P.L., see Tooyama, I., 155
Mefford, I.N., see Ota, M., 129
Mewett, K., see Fletcher, E., 133
Migheli, A., Attanasio, A., Vigliani, M.C. and Schiffer, D., Dystrophic neurites around amyloid plaques of human patients with Gerstmann–Sträussler–Scheinker disease contain ubiquitinated inclusions, 55
Miller, J.A., The calibration of ^{35}S or ^{32}P with ^{14}C-labeled brain paste or ^{14}C-plastic standards for quantitative autoradiography using LKB Ultrofilm or Amersham Hyperfilm, 211
Misgeld, U., see Jarolimek, W., 227
Modena, I., see Narici, L., 51
Morita, K., see Nakanishi, A., 163
Morris, C., see Kerwin, J., 178
Mortensen, A., see Edvinsson, L., 151
Mosconi, T.M. and Rice, F.L., Sensory innervation of the mystacial pad fur of the ferret, 199
Murakumo, Y., see Nakanishi, A., 163

Näätänen, R., see Sams, M., 43
Nakamura, K., see Ono, T., 194
Nakanishi, A., Yoshizumi, M., Morita, K., Murakumo, Y., Houchi, H. and Oka, Motoo, Palytoxin: a potent stimulator of catecholamine release from cultured bovine adrenal chromaffin cells, 163
Nakayama, H., Nakashima, T. and Kurogochi, Y., α4 is a major acetylcholine binding subunit of cholinergic ligand affinity-purified nicotinic acetylcholine receptor from rat brains, 122

Itzhaki, R.F., see Guy, S.P., 166
Nakashima, T., see Nakayama, H., 122
Naoi, M., see Ota, M., 129
Narici, L., Modena, I., Opsomer, R.J., Pizzella, V., Romani, G.L., Torrioli, G., Traversa, R. and Rossini, P.M., Neuromagnetic somatosensory homunculus: a non-invasive approach in humans, 51
Neary, J., see Crow, T., 107

Oakley, A., see Kerwin, J., 178
Oka, Motoo, see Nakanishi, A., 163
Ono, T., Nakamura, K., Fukuda, M. and Tamura, R., Place recognition responses of neurons in monkey hippocampus, 194
Opsomer, R.J., see Narici, L., 51
Ota, M., Mefford, I.N., Naoi, M. and Linnoila, M., Effects of morphine administration on catecholamine levels in rat brain; specific reduction of epinephrine concentration in hypothalamus, 129

Pansky, B., see Das, A., 231
Parnavelas, J.G., see Dinopoulos, A., 25
Pelletier, J., see De Reviers, M.-M., 17
Peterson, C., Vanderklish, P., and Seubert, P., Increased spectrin proteolysis in fibroblasts from aged and Alzheimer donors, 239
Perry, E., see Kerwin, J., 178
Perry, R., see Kerwin, J., 178
Pizzella, V., see Narici, L., 51
Pockett, S. and Slack, J.R., Presynaptic activity in the period immediately after conditioning is not necessary for long-term potentiation, 81
Post, C., see Karlsten, R., 267

Reid, T.W., see Das, A., 231
Reymann, K.G., see Matthies Jr., H., 259
Rice, F.L., see Mosconi, T.M., 199
Rogers, L.J. and Bolden, S.W., Light-dependent development and asymmetry of visual projections, 63
Romani, G.L., see Narici, L., 51
Rose, D., see Larnicol, N., 93
Rossini, P.M., see Narici, L., 51

Sacaan, A.I. and Johnson, K.M., Regulation of polyamine and magnesium inhibition of the *N*-methyl-D-aspartate (NMDA) receptor ionophore complex, 219
Sams, M. and Näätänen, R., Neuromagnetic responses of the human auditory cortexto short frequency glides, 43
Sandner, G. and Scala, G.D., Periaqueductal gray spike trains recorded in frontal or horizontal mesencephalic brain slices from the rat, 147
Sato, M., Ikeda, K., Haga, S., Allsop, D. and Ishii, T., A monoclonal antibody to common acute lymphoblastic leukemia antigen (neutral endopeptidase) immunostains senile plaques in the brains of patients with Alzheimer's disease, 271
Scala, G.D., see Sandner, G., 147
Schiffer, D., see Migheli, A., 55
Schmidt-Kastner, R., see Wagemann, E., 102
Schroeder, R.L., Weinger, M.B., Vakassian, L. and Koob, G.F., Methylnaloxonium diffuses out of the rat brain more slowly than naloxone after direct intracerebral injection, 173
Schütte, M., [^{125}I]SCH 23982, a new tool for rapid visualization of dopaminergic neurons in lower vertebrate retinas, 29
Sebens, J., see Jaarsma, D., 111
Seubert, P., see Peterson, C., 239
Sherman, A.D., Davidson, A.T., Baruah, S., Hegwood, T.S. and Waziri, R., Evidence of glutamatergic deficiency in schizophrenia, 77
Shippenberg, T.S., Conditioned reinforcing effects of 8-hydroxy-2-(di-*N*-propylamino) tetralin: involvement of 5-hydroxytryptamine 1*A* and D_1 dopamine receptors, 136

Sierra, A., see Aceves, J., 223
Slack, J.R., see Pockett, S., 81
Smart, T.G., see Bowie, D., 68
Smith, P.F., Darlington, C.L. and Hubbard, J.I., Evidence for inhibitory amino acid receptors on guinea pig medial vestibular nucleus neurons in vitro, 244
So, K. -F., see Lau, K.C., 255
Sontag, K.-H., see Wagemann, E., 102
Speckmann, E. -J., see Lehmenkühler, C., 5
Sugiyama, H., see Ito, I., 119
Susel, Z., Engber, T.M., Kuo, S. and Chase, T.N., Prolonged infusion of quinolinic acid into rat striatum as an excitotoxic model of neurodegenerative disease, 234

Takenaka, T., see Horie, H., 125
Tamura, R., see Ono, T., 194
Tanaka, K., Koyano, K. and Kuba, K., A muscarine-activated voltage-independent K channel in cultured bullfrog sympathetic neurones, 191
Tay, D., see Lau, K.C., 255
Ter Horst, G.J., see Copray, J.C.V.M., 97
Terakado, Y., see Ichikawa, Y., 37
Tierney, B.E., see Keller, J.T., 247
Tillet, Y., see De Reviers, M.-M., 17
Tooyama, I., Akiyama, H., McGeer, P.L., Hara, Y. Yasuhara, O. and Kimura, H., Acidic fibroblast growth factor-like immunoreactivity in brain of Alzheimer patients, 155
Torrioli, G., see Narici, L., 51
Traversa, R., see Narici, L., 51
Treanor, J.J.S., Dawbarn, D., Allen, S.J., MacGowan, S.H. and Wilcock, G.K., Low affinity nerve growth factor receptor binding in normal and Alzheimer's disease basal forebrain, 73

Uddman, R., see Edvinsson, L., 151

Vaillier, J., see Chapron, J., 21
Vakassian, L., see Schroeder, R.L., 173
Van Willigen, J.D., see Copray, J.C.V.M., 97
Vanderklish, P., see Peterson, C., 239
Veh, R.W., Digoxigenylated wheat germ agglutinin visualized with alkaline phosphatase-labeled anti-digoxigenin antibodies – a new, sensitive technique with the potential for single and double tracing of neuronal connections, 169
Vigliani, M.C., see Migheli, A., 55

Wagemann, E., Schmidt-Kastner, R., Block, F. and Sontag, K.-H., Neuronal degeneration in hippocampus and cerebellum of mutant spastic Han-Wistar rats, 102
Walden, J., see Lehmenkühler, C., 5
Wander, A.H., see Keller, J.T., 247
Waxham, M.N., see Crow, T., 107
Waziri, R., see Sherman, A.D., 77
Weight, F.F., see Aguayo, L.G., 88
Weinger, M.B., see Schroeder, R.L., 173

Westrum, L.E. and Henry, M.A., Contralateral degeneration in the cat spinal trigeminal nucleus following unilateral retrogasserian trigeminal rhizotomy, 143
White, G., see Aguayo, L.G., 88
Wilcock, G.K., see Treanor, J.J.S., 73
Williams, M., see Crow, T., 107
Williamson, R. and Budelmann, B.U., Convergent inputs to *Octopus* oculomotor neurones demonstrated in a brain slice preparation, 215
Wood, J.G. and Zinsmeister, P., Tyrosine phosphorylation systems in Alzheimer's disease pathology, 12

Yamada, K., see Kohmura, E., 159
Yamaguchi, T., see Blin, J., 183
Yamaguchi, T., see Ichikawa, Y., 37
Yamasaki, Y., Kogure, K., Hara, H., Ban, H. and Akaike, N., The possible involvement of tetrodotoxin-sensitive ion channels in ischemic neuronal damage in the rat hippocampus, 251
Yasuhara, O., see Tooyama, I., 155
Yoshizumi, M., see Nakanishi, A., 163

Zagami, A.S. and Lambert, G.A., Craniovascular application of capsaicin activates nociceptive thalamic neurones in the cat, 187
Zinsmeister, P., see Wood, J.G., 12
Zvolský, P., see Burešová, M., 47

ANNOUNCEMENT

The *17th International Summer School of Brain Research on the Human Hypothalamus in Health and Disease* will be held in Amsterdam, The Netherlands, on August 26–30, 1991.

The aim of the Summer School is to provide high-level didactic reviews on the state of affairs in the research on the human hypothalamus. Topics that will be discussed are normal development, sexual differentiation and aging of the human hypothalamus, and the pathophysiological alterations in neurological disease.

The Summer School intends to integrate, for the first time, the recent data on the human hypothalamus provided by various international groups from different disciplines for students, researchers and clinicians.

For further information: Ms. Tini Eikelboom or Ms. Olga Pach
Netherlands Institute for Brain Research
Meibergdreef 33
1105 AZ Amsterdam ZO
The Netherlands
Tel. 20 - 5665500
Fax 20 - 918466